

AMENDMENTS TO THE CLAIMS

1. (Currently amended) A high-throughput method for producing a plurality of monoclonal ~~antibody~~ antibodies, each of which binds to a different candidate antigen, said method comprising the steps of:

- a) introducing ~~at least one~~ a plurality of candidate ~~antigen~~ antigens into an animal or animals;
- b) recovering antibody-producing cells from said animal or animals and rendering these cells into a single cell suspension;
- c) generating ~~an~~ immortalized cell ~~line~~ lines from said single cell suspension;
- d) screening the supernatant of said immortalized cell ~~line~~ lines against a protein chip or protein chips on which the candidate ~~antigen is~~ antigens are displayed; and
- (e) selecting ~~as said monoclonal antibody, an antibody~~ antibodies that ~~binds~~ bind to said candidate ~~antigen~~ antigens.

2. (Currently amended) The method of claim 1, wherein said animal or animals is a mouse, a rat, a guinea pig or a rabbit.

3. (Cancelled)

4. (Currently amended) The method of claim 3 1, wherein between ~~one~~ two and fifty different purified candidate antigens are introduced into ~~the~~ each animal.

5. (Currently amended) The method of claim 4, wherein between 0.001 and 1000 micrograms of each antigen is introduced into ~~the~~ each animal.

6. (Currently amended) The method of claim 1 comprising the additional step of supplying the animal or animals with a booster dose of some or all of the antigens which were introduced into the animal or animals prior to the removal of antibody-producing cells.

7. (Currently amended) The method of claim 1, wherein the antibody-producing cells are B cells, T cell or stem cells.

8. (Currently amended) The method of claim 1, wherein the antibody-producing cells are recovered by removal of spleen tissue, lymph nodes or bone marrow of the animal or animals.

9. (Currently amended) The method of claim 1, wherein the immortalized cell ~~line is a~~ lines are hybridoma cell ~~line~~ lines produced by somatic fusion of the cells in the single cell suspension to myeloma cells.

10. (Currently amended) The method of claim 1, wherein said protein chip or protein chips is a plain-glass slide, a 3D gel pad chip, a microwell chip or a cell chip.

11. (Currently amended) The method of claim 1, wherein the step of detecting the monoclonal antibodies bound to the antigens further comprises isotyping the monoclonal antibodies.

12. (Currently amended) The method of claim 11, wherein said step of detecting and isotyping the monoclonal antibodies comprises adding isotype specific anti-immunoglobulin antibodies to said protein chip or protein chips, wherein each anti-immunoglobulin antibody having a different isotype specificity has a different label, and detecting the presence of said labels.

13. (Currently amended) The method of claim 1, further comprising assessing the specificity with which each isolated monoclonal antibody binds to an antigen using a protein chip or protein chips comprising said antigen.

14-15. (Cancelled)

16. (Withdrawn) A method for producing an immortalized cell line that produces a monoclonal antibody of interest, said method comprising the steps of:

- a) introducing at least one candidate antigen into an animal;
- b) recovering antibody-producing cells from said animal and rendering these cells into a single cell suspension;
- c) generating an immortalized cell line from said single cell suspension;
- d) screening the supernatant of said immortalized cell line against a protein chip on which the candidate antigen is displayed; and
- e) selecting as said immortalized cell line, that which produces a supernatant containing an antibody that binds to said candidate antigen.

17. (Withdrawn) An immortalized cell line isolated by the method of claim 16.

18. (Currently amended) A method for producing a plurality of monoclonal antibodies, each of which binds to a different purified candidate antigen, comprising introducing a plurality of purified candidate antigens into an animal, each purified candidate antigen being ~~derived~~ obtained from a different ~~source~~ protein.

19. (Currently amended) A method for producing a plurality of monoclonal antibodies, each of which binds to a different purified candidate antigen, comprising introducing a plurality of purified candidate antigens into an animal, each purified candidate antigen being ~~derived~~ obtained from a different ~~source~~ protein, which further comprises any of the steps recited in claim 1.

20. (Withdrawn) A monoclonal antibody isolated by the method of claim 1.

21. (Withdrawn) An antibody according to claim 20 which is an anti-idiotypic antibody.

22. (Withdrawn) An antibody according to claim 21 which is an anti-anti-idiotypic antibody.

23. (Withdrawn) An immortalized cell line producing a monoclonal antibody of claim 20.

24. (Withdrawn) An immortalized cell line according to claim 23 which is a hybridoma cell line.

25. (Withdrawn) A bank of antibodies according to claim 20.

26. (Withdrawn) A bank of immortalized cell lines according to claim 15.

27. (New) A method of identifying a plurality of monoclonal antibodies, each of which binds to a different candidate antigen, said method comprising the steps of:

- a) screening the supernatant of immortalized cell lines against one or more protein chips on which the candidate antigens are displayed; and
- b) selecting monoclonal antibodies that bind to said candidate antigens, said method being characterized in that said immortalized cell lines are generated from a single suspension of antibody-producing cells that produce antibodies against a plurality of antigens.